

## Decay of $^{200,201}\text{Fr}^*$

Z. Kalaninová<sup>†1</sup>, S. Antalic<sup>1</sup>, A.N. Andreyev<sup>2,3</sup>, F.P. Heßberger<sup>4,5</sup>, D. Ackermann<sup>4</sup>, B. Andel<sup>1</sup>, L. Bianco<sup>6</sup>, S. Hofmann<sup>4</sup>, M. Huyse<sup>7</sup>, B. Kindler<sup>4</sup>, B. Lommel<sup>4</sup>, R. Mann<sup>4</sup>, R.D. Page<sup>6</sup>, P. Sappale<sup>6</sup>, J. Thomson<sup>6</sup>, P. Van Duppen<sup>7</sup>, and M. Venhart<sup>8,1</sup>

<sup>1</sup>Comenius University, Bratislava, Slovakia; <sup>2</sup>University of York, York, UK; <sup>3</sup>ASRC, JAEA, Ibaraki, Japan; <sup>4</sup>GSI, Darmstadt, Germany; <sup>5</sup>Helmholtz Institut Mainz, Mainz, Germany; <sup>6</sup>University of Liverpool, Liverpool, UK; <sup>7</sup>KU Leuven, Leuven, Belgium; <sup>8</sup>Institute of Physics, SAS, Bratislava, Slovakia

In the region of neutron-deficient nuclei above lead several interesting nuclear-structure phenomena can be observed, e.g., coexistence of states with different shapes within one nucleus, or  $\beta$ -delayed fission. This motivated us to investigate the neutron-deficient isotopes  $^{200,201}\text{Fr}$ .

The studied nuclei were produced at the velocity filter SHIP (GSI, Darmstadt) in fusion-evaporation reactions  $^{56}\text{Fe} + ^{147,149}\text{Sm}$  at several beam energies from 236 to 275 MeV. Evaporation residues (ERs) were separated from other particles and transported into a focal-plane detector system. ERs were implanted into a 16-strip position-sensitive silicon detector (PSSD) recording also their  $\alpha$  decays. Escaping  $\alpha$  particles were recorded by a system of six silicon detectors placed upstream the beam covering 80 % of  $2\pi$ . A germanium clover detector placed closely behind the PSSD registered  $\gamma$  and X-rays.

We measured  $E_\alpha = 7470(5)$  keV and  $T_{1/2} = 46(4)$  ms for  $^{200}\text{Fr}$  confirming known  $\alpha$ -decay data for this isotope. For its daughter isotope,  $^{196}\text{At}$ , we observed a new weak  $\alpha$  line at 6732(8) keV with a relative intensity of 4(2)% besides the main 7045(5)-keV  $\alpha$  line. The determined energy of the level in  $^{192}\text{Bi}$  populated by the 6732(8)-keV decay is 320(10) keV. Within a 5- $\mu\text{s}$  coincidence time with implanted ERs followed by  $\alpha$  decays of  $^{200}\text{Fr}$  we observed weak  $\gamma$  lines at 75.5 and 77.1 keV and  $K_\alpha(\text{Fr})$  X-rays. They indicate a short-lived  $\gamma$ -decaying state in  $^{200}\text{Fr}$  with  $T_{1/2} = 0.6_{-0.2}^{+0.5}$   $\mu\text{s}$ . One  $\beta$ -delayed fission ( $\beta\text{DF}$ ) event attributed to  $^{200}\text{Fr}$  was observed. Deduced probability of  $\beta\text{DF}$  for the daughter isotope  $^{200}\text{Rn}$  is more than 1.4 %.

We identified a short-lived  $\gamma$ -decaying activity with  $T_{1/2} = 0.7_{-0.2}^{+0.5}$   $\mu\text{s}$  also in  $^{201}\text{Fr}$  based on the registration of  $\gamma$  and  $K_\alpha$  X-rays. From the analysis of K-shell internal conversion coefficients ( $\alpha_K$ ) [1] and estimated single-particle half-lives ( $T_{1/2,SP}$ ) according to Weisskopf [2] we suppose that observed  $\gamma$  and X-rays arise from an internal transition of  $M2$  multipolarity. We tentatively assigned the spin and parity of  $13/2^+$  to the observed isomeric state in  $^{201}\text{Fr}$ . The lower energy limit for this level was determined to be higher than the K-shell atomic-electron binding energy of francium (101.13 keV) because of the detection of K X-rays. The upper energy limit was roughly estimated to be 300 keV from the comparison of experimental and expected  $\alpha_K$  and  $T_{1/2,SP}$  for  $M2$  transitions.

For most of the neutron-deficient francium ( $Z = 87$ )

\* Work supported by Slovak Research and Development Agency and Slovak grant agency VEGA.

<sup>†</sup> Zdenka.Kalaninova@fmph.uniba.sk




Figure 1: (Color online) Energy level systematics for odd- $A$  astatine (dashed lines and open symbols) and francium (solid lines and full symbols) isotopes.

and astatine ( $Z = 85$ ) isotopes a  $9/2^-$  state related to a spherical shape was identified to be a ground state. In astatine isotopes, the  $7/2^-$ ,  $1/2^+$ , and  $13/2^+$  levels, related to oblate shapes, were observed with energies decreasing at decreasing  $N$  (see Fig 1). Starting with  $^{195}\text{At}$  ( $N = 110$ ), the  $1/2^+$  level becomes the ground state in astatine isotopes [3]. The energy interval of the tentative  $13/2^+$  level in  $^{201}\text{Fr}$  estimated from our data follows the trend of decreasing energies at decreasing  $N$  of this level in francium isotopes. A similar trend was also observed for the  $1/2^+$  level. In the lightest francium isotopes we can expect a change of spin of the ground state, but it was not definitely identified so far. All of the  $13/2^+$ ,  $7/2^-$ ,  $1/2^+$  levels were reported to be detected in  $^{199}\text{Fr}$  within 300 keV [4]. However, in recent measurements at SHIP we only observed the  $7/2^-$  level, and tentatively also the  $1/2^+$  level [5]. Higher statistics are needed to disentangle the level structure in this isotope.

## References

- [1] T. Kibédi, *et al.*, *Nucl. Instr. and Meth. A* **589**, 202 (2008).
- [2] V.F. Weisskopf, *Phys. Rev.* **83**, 1073 (1951).
- [3] H. Kettunen, *et al.*, *Eur. Phys. J. A* **16**, 457 (2003).
- [4] J. Uusitalo, *et al.*, *Phys. Rev. C* **87**, 064304 (2013).
- [5] Z. Kalaninová, *et al.*, *Phys. Rev. C* **87**, 044335 (2013).