

Observation of $\eta_c(1S,2S)$ and χ_{cJ} decays to $2(\pi^+\pi^-)\eta$ via $\psi(3686)$ radiative transitions

M. Ablikim *et al.*^{*}
(BESIII Collaboration)

 (Received 13 June 2024; accepted 5 February 2025; published 31 March 2025)

Based on $(2712.4 \pm 14.1) \times 10^6$ $\psi(3686)$ decays collected with the BESIII detector, we have observed, for the first time, the hadronic decays of S- and P-wave charmonium states into $2(\pi^+\pi^-)\eta$ via radiative transitions from $\psi(3686)$. The branching fraction of the decay $\eta_c(1S) \rightarrow 2(\pi^+\pi^-)\eta$ has a significant dependence on the interference pattern between $\eta_c(1S)$ and non- $\eta_c(1S)$ processes. We measure it in both the destructive and constructive interference scenarios for the first time. The mass and width of the $\eta_c(1S)$ are measured to be $M = (2984.14 \pm 0.13 \pm 0.38)$ MeV/ c^2 and $\Gamma = (28.82 \pm 0.11 \pm 0.82)$ MeV, respectively. Clear signals for the decays of the $\chi_{cJ}(J = 0, 1, 2)$ and the $\eta_c(2S)$ to $2(\pi^+\pi^-)\eta$ are also observed for the first time, and the corresponding branching fractions are measured. The ratio of the branching fractions between the $\eta_c(2S)$ and $\eta_c(1S)$ decays is significantly lower than the theoretical prediction, which might suggest different dynamics in their decays.

DOI: [10.1103/PhysRevD.111.052013](https://doi.org/10.1103/PhysRevD.111.052013)

I. INTRODUCTION

Our understanding of the charmonium states with masses below the open-charm production threshold is far from satisfactory. This is especially true for the spin singlet states [1], which include the S-wave ground state η_c ($\eta_c \equiv \eta_c(1S)$) and its first radial excitation $\eta_c(2S)$. The $\eta_c(2S)$ state was first observed by the Belle experiment [2] and later confirmed by the BABAR [3] and CLEO [4] experiments. BESIII was the first to observe the radiative transition $\psi(3686) \rightarrow \gamma\eta_c(2S)$ [5]. However, we have a poor knowledge about the decays of $\eta_c(2S)$, even twenty years after its first discovery. Only about 6% of the $\eta_c(2S)$ decays have been observed and measured. By contrast, people have more studies on η_c . However, there are still many unknown decay modes, also many measured BF's have large uncertainties due to the size of data sample, background level, imperfect understanding to its mass shape, ignoring the interference between η_c and nonresonance, and so on.

The measurement of the branching fractions (BFs) of η_c and $\eta_c(2S)$ is crucial for understanding the nature of charmonium states. In the potential model, the ratio of branching fractions of J/ψ and $\psi(3686)$ decaying into the same light hadron final states is around 12% [6]. This rule

has been confirmed in various modes [7]; however, it was soon discovered to be violated in the decays into pseudo-scalar vector pairs and vector tensor pairs, which are suppressed by at least an order of magnitude. The expected ratio of BF's for decays of the η_c and $\eta_c(2S)$ into the same hadronic final states (h) was initially anticipated to be similar to the ratio (12%) observed for their spin-triplet partners J/ψ and $\psi(3686)$ [6,8].

$$\frac{\mathcal{B}(\eta_c(2S) \rightarrow h)}{\mathcal{B}(\eta_c \rightarrow h)} \simeq \frac{\mathcal{B}(\psi(3686) \rightarrow h)}{\mathcal{B}(J/\psi \rightarrow h)} = 0.128. \quad (1)$$

However, a more in-depth analysis suggested that this ratio should actually be close to unity, as the dominant decay dynamics of the η_c and $\eta_c(2S)$ involve the annihilation of charmed quark pairs into two gluons [9]. Experimental measurements have revealed inconsistencies with both of these theoretical predictions [10]. Further measurements of additional decay modes of the η_c and $\eta_c(2S)$, in addition to testing these predictions, may provide insight into the internal structure of these states, for example by testing for the presence of noncharmonium components in their wave functions.

Previous measurements of the η_c mass and width, obtained through the magnetic dipole (M1) transition $\psi(3686) \rightarrow \gamma\eta_c$, $p\bar{p}$ collisions, or $\gamma\gamma$ collisions, have shown significant discrepancies between different experiments [11], which are possibly caused by many reasons mentioned before. However, interference between η_c decays and nonresonant processes could also have a large impact on these measurements as well as on the η_c BF's. The CLEO [12], BESIII [13], and KEDR [14] collaborations have studied η_c decays using

^{*}Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

the M1 transition $\psi(3686) \rightarrow \gamma\eta_c$, which is considered to be a golden channel for η_c production due to its low background level and the few unknown background channels. BESIII successfully described the η_c line shape by using the energy dependence of the hindered-M1 transition matrix element and by fully considering interference with nonresonant $\psi(3686)$ radiative decays, leading to a significant improvement in the precision of the η_c mass and width measurements [13]. However, further improvements are still beneficial since, for example, the determination of the hyperfine mass splitting of S-wave charmonium can test theoretical calculations [15,16], and the width directly relates to the theory of η_c -glueball mixing [17].

As for the P -wave spin-triplet states $\chi_{c1,2}$, the dominant decay modes are the transitions $\chi_{c1,2} \rightarrow \gamma J/\psi$. However, the sum of all measured $\chi_{c0,1,2}$ BFs are each still far less than 1. Intensive studies of their multibody decays are lacking relative to their few-body decays. The search for more new decay modes of the $\chi_{c0,1,2}$ is useful in understanding their properties.

In this article, we analyze the hadronic decays of S- and P -wave charmonium states into $2(\pi^+\pi^-)\eta$ with $(2712.4 \pm 14.1) \times 10^6$ $\psi(3686)$ events [18]. The process of $\eta_c(2S)$ and χ_{cJ} decays to $2(\pi^+\pi^-)\eta$ are first observed. With this largest $\psi(3686)$ sample in the world, better knowledge of the background and the interference between the η_c and non- η_c components, we improve the measurements of the η_c mass and width, as well as the BF of $\eta_c \rightarrow 2(\pi^+\pi^-)\eta$. Additionally, we present the first measurements of the ratio of branching fractions for η_c and $\eta_c(2S)$ decays into $2(\pi^+\pi^-)\eta$ and compare it with the theoretical calculations.

II. BESIII DETECTOR AND MONTE CARLO SIMULATION

The BESIII detector is a magnetic spectrometer [19,20] operating at the BEPCII collider [21]. The cylindrical core of the BESIII detector covers 93% of the 4π solid angle and consists of a helium-based multilayer drift chamber (MDC), a plastic scintillator time-of-flight system (TOF), and a CsI (TI) electromagnetic calorimeter (EMC), which are all enclosed in a superconducting solenoidal magnet, providing a 1.0 T magnetic field. The solenoid is supported by an octagonal flux-return yoke, with resistive plate chamber muon identifier modules interleaved with steel. The charged-particle momentum resolution at 1 GeV/ c is 0.5%, and the dE/dx resolution is 6% for the electrons from Bhabha scattering. The EMC measures photon energies with a resolution of 2.5% (5%) at 1 GeV in the barrel (end cap) region. The time resolution of the TOF barrel section is 68 ps, while that of the end cap section is 110 ps. The end cap TOF system was upgraded in 2015 with multigap resistive plate chamber technology, providing a time resolution of 60 ps [22–24].

Simulated Monte Carlo (MC) samples are produced with the GEANT4-based [25] software, which models the

experimental conditions, including the electron-positron collision, the decays of the particles, and the response of the detector. The simulation models the beam energy spread and initial state radiation (ISR) in the e^+e^- annihilations with the generator KKMC [26]. All particle decays are modeled with EVT GEN [27,28] using BFs either taken from the Particle Data Group (PDG) [11], when available, or otherwise estimated with LUNDCHARM [29,30]. Final-state radiation (FSR) from charged final-state particles is incorporated using the PHOTOS package [31]. The exclusive decays of $\psi(3686) \rightarrow \gamma X$ are generated following the angular distribution of $(1 + \lambda \cos^2 \theta_\gamma)$, where X refers to $\eta_c(1S, 2S)$ or χ_{cJ} , θ_γ is the polar angle of the radiative photon in the rest frame of the $\psi(3686)$, and the value of λ is set to 1 for $\eta_c(1S, 2S)$ and to 1, $-1/3$, $1/13$ for χ_{c0} , χ_{c1} , and χ_{c2} respectively [13,32]. The $X \rightarrow 2(\pi^+\pi^-)\eta$ decays are generated uniformly in phase space (PHSP).

III. EVENT SELECTION

Charged tracks detected in the multilayer drift chamber (MDC) are required to be within the polar angle (θ) range of $|\cos \theta| < 0.93$. Here θ is defined with respect to the z -axis, which is the symmetry axis of the MDC. For each charged track, the distance of closest approach to the interaction point must be less than 10 cm along the z -axis and less than 1 cm in the transverse plane. Photon candidates are reconstructed from isolated showers in the electromagnetic calorimeter (EMC). The deposited energy of each shower is required to be at least 25 MeV in the barrel region ($|\cos \theta| < 0.80$) and 50 MeV in the end cap region ($0.86 < |\cos \theta| < 0.92$). For the $\eta_c(2S)/\chi_{cJ}$ modes, the deposited energy is required to be larger than 40 MeV in the end cap region due to the low energy of the M1 transition photon. To exclude showers originating from charged tracks, the angle between the shower direction and the charged tracks extrapolated to the EMC must be greater than 10° . To suppress electronic noise and energy depositions unrelated to the event, the EMC cluster timing from the reconstructed event start time is required to be within [0, 700] ns.

Candidate events are required to have four charged tracks with zero net charge and at least three photons. All the charged tracks are taken as pions. To reconstruct the $\eta \rightarrow \gamma\gamma$ decay, the invariant mass of a pair of photons is required to satisfy $M_{\gamma\gamma} \in [0.51, 0.57]$ GeV/ c^2 . Then a one-constraint (1C) kinematic fit is performed for each combination with $\chi^2_{1C}(\gamma\gamma) < 20$, where the photon pair is constrained to the η mass, and at least one η candidate is required. To suppress the background and improve the mass resolution, a five-constraint (5C) kinematic fit is performed to the total initial four momentum of the colliding beams. The extra 1C is used to constrain the $M_{\gamma\gamma}$ to the η nominal mass. If there are multiple combinations in an event, the one with the minimum χ^2 value of the 5C kinematic fit (χ^2_{5C}) is chosen. Furthermore, the criterion $\chi^2_{5C} < 15$ is required, which is

determined by optimizing the figure of merit, defined as $S/\sqrt{S+B}$, where $S(B)$ is the number of signal (background) events in the fit region. The number of background events is estimated using the inclusive MC sample with the same selection criteria as for data. In addition, to suppress backgrounds from processes with missing or additional photons, a four-constraint (4C) kinematic fit is performed for both the signal and background channels. For the $\gamma\eta_c$ mode, the $\chi_{4C}^2(3\gamma 2(\pi^+\pi^-))$ is required to be less than $\chi_{4C}^2(4\gamma 2(\pi^+\pi^-))$ and $\chi_{4C}^2(2\gamma 2(\pi^+\pi^-))$. For the $\gamma\eta_c(2S)/\chi_{cJ}$ modes, only $\chi_{4C}^2(3\gamma 2(\pi^+\pi^-)) < \chi_{4C}^2(4\gamma 2(\pi^+\pi^-))$ is required since the events with one missing photon mainly come from FSR, which will be discussed later. Backgrounds from the continuum process $e^+e^- \rightarrow q\bar{q}$ are investigated using the off-resonance data sample taken at the center-of-mass (CM) energy of 3.65 GeV. It is found that its shape is generally smooth in the $\eta_c(1S, 2S)$ and χ_{cJ} mass regions.

IV. BACKGROUND ESTIMATION

Background events from the $\psi(3686) \rightarrow PJ/\psi(P = \pi^+\pi^-, \eta)$ process are suppressed by requiring the recoil mass of P to not fall in the J/ψ signal region, approximately (3.090, 3.104) GeV/ c^2 . In case of multiple $\pi^+\pi^-$ combinations in an event, we select the one with M_P^{recoil} closest to the J/ψ mass. Background events from the $\psi(3686) \rightarrow \pi^0 H$ process (H denotes hadronic states) are suppressed by requiring the invariant mass of all possible photon pair combinations greater than 0.15 GeV/ c^2 or less than 0.12 GeV/ c^2 . For $\gamma\eta_c(2S)/\chi_{cJ}$ decay modes, the number of background events from the $\psi(3686) \rightarrow \gamma\chi_{cJ}$; $\chi_{cJ} \rightarrow \gamma J/\psi$ process is suppressed by requiring the recoil invariant mass of all possible photon pair combinations greater than 3.137 GeV/ c^2 or less than 3.052 GeV/ c^2 . For $\psi(3686) \rightarrow \gamma\eta_c$ decay mode, the fake η background due to the photon wrong combination is found. It is suppressed by requiring the invariant mass of the wrong-combination photon pair greater than 0.57 GeV/ c^2 or less than 0.53 GeV/ c^2 .

A. $\psi(3686) \rightarrow 2(\pi^+\pi^-)\eta$

The process $\psi(3686) \rightarrow 2(\pi^+\pi^-)\eta$ can contaminate our signal channel through FSR from one of the final-state pions. This effect is particularly notable in the $\eta_c(2S)$ mass region, where the FSR photon has a similar energy to the radiative photon. To separate this background from the $\eta_c(2S)$ signal, we perform a modified kinematic fit where the energy of the radiative photon is floating in the 5C fit (m5C). Although the resolution of the $\eta_c(2S)$ signal itself becomes slightly worse, the FSR background shape becomes smoother. This results in a widened gap between the $\eta_c(2S)$ peak and FSR background [32]. Furthermore, the background shape is sensitive to the fraction of the number of background events between FSR and non-FSR

TABLE I. Estimates of potential peaking backgrounds.

Background	χ_{c0}	χ_{c1}	χ_{c2}
$\chi_{c0} \rightarrow \eta'\eta'; \eta' \rightarrow \gamma\pi^+\pi^-$	288 ± 17	0	0
$\chi_{cJ} \rightarrow \gamma J/\psi; J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$	0	235 ± 21	113 ± 11
$\chi_{cJ} \rightarrow \gamma J/\psi; J/\psi \rightarrow \gamma 2(\pi^+\pi^-)$	0	55 ± 10	2.7 ± 1.7

modes. The discrepancy in the FSR fraction between data and MC simulation is studied using a control sample $\psi(3686) \rightarrow \gamma\chi_{c0}, \chi_{c0} \rightarrow 3(\pi^+\pi^-)(\gamma_{\text{FSR}})$, and subsequently corrected by the FSR fraction. The FSR photon in the control sample is required to be reconstructed, and the event selection criteria for the control sample are similar to those described in Ref. [32]. The background shape from $\psi(3686) \rightarrow 2(\pi^+\pi^-)\eta$ will be described by the sum of MC simulated shapes $\psi(3686) \rightarrow 2(\pi^+\pi^-)\eta$ and $\psi(3686) \rightarrow 2(\pi^+\pi^-)\eta\gamma_{\text{FSR}}$ with the corrected FSR fraction.

B. $\psi(3686) \rightarrow \pi^0 2(\pi^+\pi^-)\eta$

The process $\psi(3686) \rightarrow \pi^0 2(\pi^+\pi^-)\eta$ is the dominant background where a photon from the π^0 decay is missing, and it is more serious in the η_c mass region. A background events reconstruction method [13] is employed to determine its contribution by comparing the detection efficiencies between the signal and background channels, and combining with the actual number of events observed in data. The yields for the $\psi(3686) \rightarrow \pi^0 2(\pi^+\pi^-)\eta$ process in each $M(2(\pi^+\pi^-)\eta)$ bin are extracted by fitting to the π^0 signal in $M(\gamma\gamma)$ spectrum. The contribution of $\psi(3686) \rightarrow \pi^0 2(\pi^+\pi^-)\eta$ will be fixed according to the data-driven result when fitting in the η_c mass region.

C. Estimation of peaking backgrounds

After applying the same event selection criteria to the inclusive $\psi(3686)$ MC events, which are almost the same size as for data, an event-type analysis tool TopoAna [33] is used to analyse the background. We find that there are several potential peaking backgrounds for the χ_{cJ} modes. The normalized contributions for these backgrounds are listed in Table I, using branching fractions from the PDG [11] and efficiencies from the simulated exclusive MC samples. We will directly subtract these contributions from the number of signal events when calculating the BF.

V. SIGNAL YIELDS

A. $\chi_{cJ}/\eta_c(2S) \rightarrow 2(\pi^+\pi^-)\eta$

The signal yields for $\chi_{cJ}/\eta_c(2S)$ are obtained by performing an unbinned maximum likelihood fit to the $M_{2(\pi^+\pi^-)\eta}^{m5C}$ distribution, as shown in Fig. 1. The fit function includes the $\chi_{cJ}/\eta_c(2S)$ signals, continuum background, FSR contribution from $\psi(3686) \rightarrow 2(\pi^+\pi^-)\eta$, and other background contributions from $\psi(3686)$ decays. The χ_{cJ}




FIG. 1. Fit to the $M^{5C}_{2(\pi^+\pi^-\eta)}$ spectrum. The different components are expressed with different line styles or colors as given in the legend.

signal functions are described by MC-simulated shapes convolved with Gaussian functions to account for the difference in detector resolution between data and MC simulation, and the parameters of the smeared Gaussian functions are free. The signal function of $\eta_c(2S)$ is described by the Eq. (2), which has been applied in the relevant BESIII analysis [32,34–36].

$$\epsilon(m)(E_\gamma^3 \times BW(m) \times f_d(E_\gamma) \otimes DG) \otimes G(\delta m, \sigma). \quad (2)$$

where $\epsilon(m)$ is the energy-dependent detection efficiency determined by MC simulation with a PHSP model, $E_\gamma \equiv (m_{\psi(3686)}^2 - m^2)/2m_{\psi(3686)}$ is the energy of the radiative photon, and $BW(m)$ is the nonrelativistic Breit-Wigner (BW) function, in which the $\eta_c(2S)$ mass and width are fixed to those in the PDG [11]. The function $f_d(E_\gamma)$ is used to damp the diverging tail caused by the E_γ^3 term, and is given by $[E_0^2/(E_\gamma E_0 + (E_\gamma - E_0)^2)]$, as introduced by the KEDR experiment [37]. Here, $E_0 = (m_{\psi(3686)}^2 - m_{\eta_c(2S)}^2)/2m_{\psi(3686)}$ denotes the peaking energy of the transition photon; $\otimes$ is the convolution operator; DG is a double Gaussian function accounting for the detector resolution, with two parameters obtained by comparing the difference in $M_{2(\pi^+\pi^-\eta)}$ before and after going through the detector; and $G(\delta m, \sigma)$ is a Gaussian function accounting for the difference in detector resolution between data and MC simulation, where σ in the $\eta_c(2S)$ mass region is fixed to the value obtained from the linear extrapolation from the χ_{cJ} mass region, while δm is allowed to be free.

The shape of the contribution from FSR background is fixed to that obtained from the corrected MC-simulated shape, and the number of events is allowed to be free. The contribution from the continuum background is fixed to that obtained from the normalized off-resonance data with the luminosity, and shifted according to the CM energy.

TABLE II. The signal yields, detection efficiencies, and the obtained BF's for $\chi_{cJ}/\eta_c(2S) \rightarrow 2(\pi^+\pi^-\eta)$ decays. The first and second uncertainties are statistical and systematic, respectively.

Decay	N^{obs}	$\epsilon(\%)$	$\mathcal{B} (\times 10^{-3})$
$\eta_c(2S) \rightarrow 2(\pi^+\pi^-\eta)$	565 ± 30	6.27	$12.1 \pm 0.6 \pm 6.2$
$\chi_{c0} \rightarrow 2(\pi^+\pi^-\eta)$	42530 ± 240	8.90	$4.55 \pm 0.03 \pm 0.40$
$\chi_{c1} \rightarrow 2(\pi^+\pi^-\eta)$	78440 ± 290	9.31	$8.06 \pm 0.03 \pm 0.77$
$\chi_{c2} \rightarrow 2(\pi^+\pi^-\eta)$	50980 ± 240	8.62	$5.91 \pm 0.03 \pm 0.57$

The remaining background distribution from $\psi(3686)$ decay is found to be smooth and described by an ARGUS function [38].

Figure 1 shows the fit result with a goodness of fit $\chi^2/\text{ndf} = 537/360$, where ndf is the number of degrees of freedom. The insert shows the $\eta_c(2S)$ mass region on a linear scale. The numerical results are summarized in Table II. The statistical significance of the $\eta_c(2S)$ signal is determined to be 8.4σ by comparing the difference in likelihoods with and without including the $\eta_c(2S)$ signal component, taking into account the difference in ndf ($\Delta\text{ndf} = 2$).

The BF's for our studied channels are calculated via

$$\mathcal{B}(X \rightarrow 2(\pi^+\pi^-\eta)) = \frac{N^{\text{obs}}}{N_{\psi(3686)}^{\text{tot}} \times \mathcal{B}_1 \times \mathcal{B}_2 \times \epsilon}, \quad (3)$$

where N^{obs} is the number of signal events, $N_{\psi(3686)}^{\text{tot}}$ is the total number of $\psi(3686)$ events, $\mathcal{B}_1$ and $\mathcal{B}_2$ are the BF's of $\psi(3686) \rightarrow \gamma X$ and $\eta \rightarrow \gamma\gamma$ [11], respectively, and ϵ is the detection efficiency after correcting the helix parameters of charged tracks [39]. Table III lists the product BF's of $\mathcal{B}(\psi(3686) \rightarrow \gamma X) \times \mathcal{B}(X \rightarrow 2(\pi^+\pi^-\eta))$.

B. $\eta_c \rightarrow 2(\pi^+\pi^-\eta)$

The η_c signal yield is obtained by performing an unbinned maximum likelihood fit to the $M_{2(\pi^+\pi^-\eta)}^{5C}$ distribution [13]. The fit function includes the η_c signal function and four background components: the background from $\psi(3686) \rightarrow \pi^0 2(\pi^+\pi^-\eta)$, the continuum background, the nonresonant background ($\psi(3686) \rightarrow \gamma 2(\pi^+\pi^-\eta)$), and other potential backgrounds from $\psi(3686)$ decays estimated using the $\psi(3686)$ inclusive MC sample. Interference between the η_c and the nonresonant process is considered. The fit function is expressed as

$$(\epsilon(m)|e^{i\phi} E_\gamma^2 BW(m) + \alpha \mathcal{N}|^2 E_\gamma^3) \otimes G + f_{\text{BKG}}, \quad (4)$$

where the first term is the η_c signal shape, which is described by the sum of a modified BW function and a coherent amplitude squared, convolved with the resolution function. Here, $\epsilon(m)$ is the mass-dependent efficiency; ϕ is the interference phase angle; E_γ is the energy of the M1




FIG. 2. The fit results for the η_c signal with destructive (a) and constructive (b) interference solutions. The different components are expressed with different line styles or colors.

transition photon; $BW(m)$ is a nonrelativistic BW function defined as $\frac{\Gamma/2}{(m-m_0)+i\Gamma/2}$, where m_0 and Γ are the mass and width of the η_c ; and α is the strength of the nonresonant component. The nonresonant PDF, denoted as $\mathcal{N}$, is

TABLE III. The product BF of $\mathcal{B}(\psi(3686) \rightarrow \gamma X) \times \mathcal{B}(X \rightarrow 2(\pi^+\pi^-)\eta)$. The first and second uncertainties are statistical and systematic, respectively.

Decay	$\mathcal{B}(\times 10^{-5})$
$\psi(3686) \rightarrow \gamma \eta_c(2S); \eta_c(2S) \rightarrow 2(\pi^+\pi^-)\eta$	$0.85 \pm 0.04 \pm 0.13$
$\psi(3686) \rightarrow \gamma \chi_{c0}; \chi_{c0} \rightarrow 2(\pi^+\pi^-)\eta$	$44.4 \pm 0.3 \pm 3.7$
$\psi(3686) \rightarrow \gamma \chi_{c1}; \chi_{c1} \rightarrow 2(\pi^+\pi^-)\eta$	$78.7 \pm 0.3 \pm 7.4$
$\psi(3686) \rightarrow \gamma \chi_{c2}; \chi_{c2} \rightarrow 2(\pi^+\pi^-)\eta$	$55.3 \pm 0.3 \pm 5.3$
$\psi(3686) \rightarrow \gamma \eta_c; \eta_c \rightarrow 2(\pi^+\pi^-)\eta$ (destructive)	$2.01 \pm 0.17 \pm 0.64$
$\psi(3686) \rightarrow \gamma \eta_c; \eta_c \rightarrow 2(\pi^+\pi^-)\eta$ (constructive)	$0.95 \pm 0.17 \pm 0.44$

described by a second order Chebychev function. The Gaussian function G accounts for the detector resolution, and f_{BKG} denotes the sum of the other three background components mentioned earlier, excluding the nonresonant background. In the fit, ϕ and α are free parameters.

The mass-dependent efficiency is determined from MC simulations of the PHSP decay $\eta_c \rightarrow 2(\pi^+\pi^-)\eta$. We use a Gaussian function to describe the discrepancy between data and MC simulation, and the parameters are determined by fitting the J/ψ peak in the control sample $\psi(3686) \rightarrow \gamma J/\psi, J/\psi \rightarrow 2(\pi^+\pi^-)\eta$.

Figure 2 shows the fit result, where the goodness of fit is $\chi^2/\text{ndf} = 164/118$. The number of η_c signal events is determined to be $N^{\text{obs}} = (2.60 \pm 0.24) \times 10^4$ (destructive), or $N^{\text{obs}} = (1.24 \pm 0.20) \times 10^4$ (constructive). Using the η_c fitted signal yield, we determine the BF of $\eta_c \rightarrow 2(\pi^+\pi^-)\eta$, listed in Table IV, as well as the mass and width of the η_c under the two different interference scenarios.

VI. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties related to the measurements for $\mathcal{B}(\eta_c(2S)/\chi_{cJ} \rightarrow 2(\pi^+\pi^-)\eta)$ include the following sources. The tracking and PID efficiencies are all assigned to be 1.0% per charged track [40], and the photon efficiency is also assigned to be 1.0% per photon [41]. The uncertainties caused by the kinematic fit are determined to be (5.5-6.9)% by comparing the difference in detection efficiency with and without correcting the helix parameters of charged tracks in MC simulation [39]. The uncertainties caused by the background veto are determined to be (0.3-6.1)% by randomly sampling multiple mass windows [42]. The uncertainties caused by the number of continuum background events are determined to be 0.23% for the $\eta_c(2S)$ signal and negligible for the χ_{cJ} signals. They are estimated by varying the number of events in the fit by one standard deviation, assuming they follow either a Poisson or binomial distribution. The uncertainties caused by MC imperfections are estimated to be (0.1-1.3)% by comparing the differences with and without correcting the momentum distributions in the MC simulation based on data. The uncertainty related to $\mathcal{B}(\eta \rightarrow \gamma\gamma)$ is taken from the PDG [11], and the uncertainty related to $\mathcal{B}(\psi(3686) \rightarrow \gamma \eta_c(2S))$ is taken from Ref. [10]. The uncertainty due to the total number of $\psi(3686)$ events is determined to be 0.5% [18]. The uncertainties due to MC statistics are determined to be

TABLE IV. The obtained mass, width, and BF of the η_c for the two interference solutions. The first and second uncertainties are statistical and systematic, respectively.

Solution	Destructive	Constructive
Mass (MeV/ c^2)	$2984.14 \pm 0.12 \pm 0.34$	$2984.13 \pm 0.13 \pm 0.38$
Width (MeV)	$28.82 \pm 0.11 \pm 0.81$	$28.81 \pm 0.11 \pm 0.82$
$\mathcal{B}$ (%)	$5.5 \pm 0.5 \pm 1.9$	$2.6 \pm 0.4 \pm 1.3$

0.3% by the same method as for the continuum background events number. The uncertainties caused by the signal yields include the following components: continuum background shape, FSR factor, mass and width of the $\eta_c(2S)$, efficiency curve, damping function, detector resolution, and other potential background from $\psi(3686)$ decays. Among them, the uncertainty due to the efficiency curve is estimated by comparing the difference in the fit results with and without including the curve. The uncertainties caused by the continuum and the potential background shapes are estimated by alternatively employing different functions. The uncertainties from other components in signal yields are estimated by varying the corresponding parameters by one standard deviation. The resulting uncertainties due to signal yields are determined to be (0.2–11.0)%. We assume all these sources are independent and take their sum in quadrature as the total systematic uncertainties, which are determined to be 15.7% (not including the uncertainty from $\mathcal{B}(\psi(3686) \rightarrow \gamma\eta_c(2S))$ measurement), 8.9%, 9.5%, and 9.7% for $\eta_c(2S)/\chi_{cJ}$, respectively.

The uncertainties related to the measurement of $\mathcal{B}(\eta_c \rightarrow 2(\pi^+\pi^-)\eta)$ have the same or similar sources as above, and are estimated with the same or similar methods. However, the same source may introduce two different uncertainties, corresponding to the constructive and destructive solutions, due to interference between the $\eta_c(1S)$ decay and the nonresonant process. Additionally, the dominant uncertainties come from the nonresonant process with different interference fraction assumptions, which are estimated to be 30.5% (constructive) or 44.1% (destructive) by mixing different fractions within one standard deviation, and the largest differences relative to the nominal results are taken as the uncertainties. Finally, the total systematic uncertainty is determined to be 34.8% (constructive) or 48.9% (destructive).

The uncertainties related to the measurements of the mass and width of the η_c are the same as those mentioned above for the η_c branching fraction measurement and are estimated with the same methods, except for the mass scale, which is estimated using fits to the χ_{cJ} peaks in the same final state. The relative uncertainties are listed in Table IV.

VII. SUMMARY

In summary, the hadronic decays of $\chi_{cJ}/\eta_c(1S, 2S)$ states have been investigated using a sample of $(2712.4 \pm 14.1) \times 10^6$ $\psi(3686)$ radiative transitions. The BF of $\eta_c \rightarrow 2(\pi^+\pi^-)\eta$ is determined for the first time under different interference scenarios. Notably, there is a significant difference between the constructive and destructive solutions. The $\chi_{cJ}/\eta_c(2S) \rightarrow 2(\pi^+\pi^-)\eta$ decays are observed for the first time, with statistical significance greater than 8σ , and the corresponding BFs are measured. We present the most precise single measurement of the η_c mass and width by assuming all radiative nonresonant events interfere with the

η_c . The relative phase for constructive or destructive interference is consistent with the earlier BESIII result [13] within 1σ , which may suggest a common phase in all modes. The physics behind this possible common phase needs to be understood. With the measured η_c mass, we obtain the hyperfine mass splitting to be $\Delta M_{hf}(1S)_{c\bar{c}} = M(J/\psi) - M(\eta_c) = 112.8 \pm 0.4 \text{ MeV}/c^2$, which is not in conflict with theoretical calculations [15,16].

Furthermore, using the BFs measured in this analysis, the ratio of BFs is

$$\frac{\mathcal{B}(\eta_c(2S) \rightarrow 2(\pi^+\pi^-)\eta)}{\mathcal{B}(\eta_c \rightarrow 2(\pi^+\pi^-)\eta)} = \begin{cases} 0.22 \pm 0.14 & (\text{destructive}) \\ 0.47 \pm 0.34 & (\text{constructive}) \end{cases}. \quad (5)$$

The results are consistent with most experimental measurements [10], but obviously disagree with the prediction in Ref. [9]. The inconsistency between the experimental measurements and the theoretical prediction on the ratio $\mathcal{B}(\eta_c(2S) \rightarrow h)/\mathcal{B}(\eta_c \rightarrow h)$ implies that mixing of the pseudoscalar glueballs to η_c or $\eta_c(2S)$ may play an important role in charmonium decays [17,43,44]. Additionally, the distinct contributions of the open-charm loop to the decays of the η_c and $\eta_c(2S)$ cannot be ignored [45].

ACKNOWLEDGMENTS

The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts No. 2020YFA0406300, No. 2020YFA0406400, No. 2023YFA1606000; National Natural Science Foundation of China (NSFC) under Contracts No. 12375071, No. 11635010, No. 11735014, No. 11935015, No. 11935016, No. 11935018, No. 12025502, No. 12035009, No. 12035013, No. 12061131003, No. 12192260, No. 12192261, No. 12192262, No. 12192263, No. 12192264, No. 12192265, No. 12221005, No. 12225509, No. 12235017, No. 12361141819, No. 12375070; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contract No. U2032108, No. U1832207; 100 Talents Program of CAS; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; German Research Foundation DFG under Contracts No. 455635585, No. FOR5327, No. GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia;

National Science Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation of Thailand under Contract No. B16F640076; Polish

National Science Centre under Contract No. 2019/35/O/ST2/02907; The Swedish Research Council; U.S. Department of Energy under Contract No. DE-FG02-05ER41374.

-
- [1] N. Brambilla *et al.*, *Eur. Phys. J. C* **71**, 1534 (2011).
 [2] S.-K. Choi *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **89**, 102001 (2002).
 [3] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **92**, 142002 (2004).
 [4] D. M. Asner *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **92**, 142001 (2004).
 [5] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **109**, 042003 (2012).
 [6] T. Appelquist and H. D. Politzer, *Phys. Rev. Lett.* **34**, 43 (1975).
 [7] S. J. Brodsky, G. P. Lepage, and S. F. Tuan, *Phys. Rev. Lett.* **59**, 621 (1987).
 [8] M. Anselmino, M. Genovese, and E. Predazzi, *Phys. Rev. D* **44**, 1597 (1991).
 [9] K. T. Chao, Y. F. Gu, and S. F. Tuan, *Commun. Theor. Phys.* **25**, 471 (1996).
 [10] H. P. Wang and C. Z. Yuan, *Chin. Phys. C* **46**, 071001 (2022).
 [11] S. Navas *et al.* (Particle Data Group), *Phys. Rev. D* **110**, 030001 (2024).
 [12] R. E. Mitchell *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **106**, 159903 (2011).
 [13] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **108**, 222002 (2012).
 [14] V. V. Anashin *et al.*, *Phys. Lett. B* **738**, 391 (2014).
 [15] T. Burch *et al.*, *Phys. Rev. D* **81**, 034508 (2010).
 [16] T. Kawanai and S. Sasaki, *Phys. Rev. D* **85**, 091503 (2012).
 [17] R. Q. Zhang *et al.*, *Phys. Lett. B* **827**, 136960 (2022).
 [18] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **48**, 093001 (2024).
 [19] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
 [20] K.-X. Huang *et al.*, *Nucl. Sci. Tech.* **33**, 142 (2022).
 [21] C. H. Yu *et al.*, *Proceedings of IPAC2016, Busan, Korea* (2016), 10.18429/JACoW-IPAC2016-TUYA01.
 [22] X. Li *et al.*, *Rad. Det. Tech. Meth.* **1**, 13 (2017).
 [23] Y. X. Guo *et al.*, *Rad. Det. Tech. Meth.* **1**, 15 (2017).
 [24] P. Cao *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **953**, 163053 (2020).
 [25] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
 [26] S. Jadach, B. F. L. Ward, and Z. Was, *Phys. Rev. D* **63**, 113009 (2001); *Comput. Phys. Commun.* **130**, 260 (2000).
 [27] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
 [28] R. G. Ping, *Chin. Phys. C* **32**, 599 (2008).
 [29] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang, and Y. S. Zhu, *Phys. Rev. D* **62**, 034003 (2000).
 [30] R. L. Yang, R. G. Ping, and H. Chen, *Chin. Phys. Lett.* **31**, 061301 (2014).
 [31] E. Barberio and Z. Was, *Comput. Phys. Commun.* **79**, 291 (1994).
 [32] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **106**, 032014 (2022).
 [33] X. Y. Zhou *et al.*, *Comput. Phys. Commun.* **258**, 107540 (2021).
 [34] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **107**, 052007 (2023).
 [35] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **109**, 042003 (2012).
 [36] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **87**, 052005 (2013).
 [37] V. V. Anashin *et al.*, *Int. J. Mod. Phys. Conf. Ser.* **02**, 188 (2011).
 [38] H. Albrecht *et al.* (ARGUS Collaboration), *Phys. Lett. B* **241**, 278 (1990).
 [39] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **87**, 012002 (2013).
 [40] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **83**, 112005 (2011).
 [41] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **81**, 052005 (2010).
 [42] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **100**, 012003 (2019).
 [43] Y. D. Tsai *et al.*, *Phys. Rev. D* **85**, 034002 (2012).
 [44] W. Qin *et al.*, *Phys. Rev. D* **97**, 096002 (2018).
 [45] Q. Wang *et al.*, *Phys. Lett. B* **711**, 364370 (2012).
-

M. Ablikim,¹ M. N. Achasov,^{4,c} P. Adlarson,⁷⁵ R. Aliberti,³⁵ A. Amoroso,^{74a,74c} M. R. An,³⁹ Q. An,^{71,58,a} Y. Bai,⁵⁷ O. Bakina,³⁶ I. Balossino,^{29a} Y. Ban,^{46,h} V. Batozskaya,^{1,44} K. Begzsuren,³² N. Berger,³⁵ M. Berlowski,⁴⁴ M. Bertani,^{28a} D. Bettoni,^{29a} F. Bianchi,^{74a,74c} E. Bianco,^{74a,74c} J. Bloms,⁶⁸ A. Bortone,^{74a,74c} I. Boyko,³⁶ R. A. Briere,⁵ A. Brueggemann,⁶⁸ H. Cai,⁷⁶ X. Cai,^{1,58} A. Calcaterra,^{28a} G. F. Cao,^{1,63} N. Cao,^{1,63} S. A. Cetin,^{62a} J. F. Chang,^{1,58} T. T. Chang,⁴³ W. L. Chang,^{1,63}

G. R. Che,⁴³ G. Chelkov,^{36,b} C. Chen,⁴³ Chao Chen,⁵⁵ G. Chen,¹ H. S. Chen,^{1,63} M. L. Chen,^{1,58,63} S. J. Chen,⁴² S. M. Chen,⁶¹ T. Chen,^{1,63} X. R. Chen,^{31,63} X. T. Chen,^{1,63} Y. B. Chen,^{1,58} Y. Q. Chen,³⁴ Z. J. Chen,^{25,i} W. S. Cheng,^{74c} S. K. Choi,^{10a} X. Chu,⁴³ G. Cibinetto,^{29a} S. C. Coen,³ F. Cossio,^{74c} J. J. Cui,⁵⁰ H. L. Dai,^{1,58} J. P. Dai,⁷⁸ A. Dbeyssi,¹⁸ R. E. de Boer,³ D. Dedovich,³⁶ Z. Y. Deng,¹ A. Denig,³⁵ I. Denysenko,³⁶ M. Destefanis,^{74a,74c} F. De Mori,^{74a,74c} B. Ding,^{66,1} X. X. Ding,^{46,h} Y. Ding,³⁴ Y. Ding,⁴⁰ J. Dong,^{1,58} L. Y. Dong,^{1,63} M. Y. Dong,^{1,58,63} X. Dong,⁷⁶ S. X. Du,⁸⁰ Z. H. Duan,⁴² P. Egorov,^{36,b} Y. L. Fan,⁷⁶ J. Fang,^{1,58} S. S. Fang,^{1,63} W. X. Fang,¹ Y. Fang,¹ R. Farinelli,^{29a} L. Fava,^{74b,74c} F. Feldbauer,³ G. Felici,^{28a} C. Q. Feng,^{71,58} J. H. Feng,⁵⁹ K. Fischer,⁶⁹ M. Fritsch,³ C. Fritzsche,⁶⁸ C. D. Fu,¹ J. L. Fu,⁶³ Y. W. Fu,¹ H. Gao,⁶³ Y. N. Gao,^{46,h} Yang Gao,^{71,58} S. Garbolino,^{74c} I. Garzia,^{29a,29b} P. T. Ge,⁷⁶ Z. W. Ge,⁴² C. Geng,⁵⁹ E. M. Gersabeck,⁶⁷ A. Gilman,⁶⁹ K. Goetzen,¹³ L. Gong,⁴⁰ W. X. Gong,^{1,58} W. Gradl,³⁵ S. Gramigna,^{29a,29b} M. Greco,^{74a,74c} M. H. Gu,^{1,58} Y. T. Gu,¹⁵ C. Y. Guan,^{1,63} Z. L. Guan,²² A. Q. Guo,^{31,63} L. B. Guo,⁴¹ R. P. Guo,⁴⁹ Y. P. Guo,^{12,g} A. Guskov,^{36,b} T. T. Han,⁵⁰ W. Y. Han,³⁹ X. Q. Hao,¹⁹ F. A. Harris,⁶⁵ K. K. He,⁵⁵ K. L. He,^{1,63} F. H. H. Heinsius,³ C. H. Heinz,³⁵ Y. K. Heng,^{1,58,63} C. Herold,⁶⁰ T. Holtmann,³ P. C. Hong,^{12,g} G. Y. Hou,^{1,63} X. T. Hou,^{1,63} Y. R. Hou,⁶³ Z. L. Hou,¹ H. M. Hu,^{1,63} J. F. Hu,^{56,j} T. Hu,^{1,58,63} Y. Hu,¹ G. S. Huang,^{71,58} K. X. Huang,⁵⁹ L. Q. Huang,^{31,63} X. T. Huang,⁵⁰ Y. P. Huang,¹ T. Hussain,⁷³ N. Hüsken,^{27,35} W. Imoehl,²⁷ M. Irshad,^{71,58} J. Jackson,²⁷ S. Jaeger,³ S. Janchiv,³² J. H. Jeong,^{10a} Q. Ji,¹ Q. P. Ji,¹⁹ X. B. Ji,^{1,63} X. L. Ji,^{1,58} Y. Y. Ji,⁵⁰ Z. K. Jia,^{71,58} P. C. Jiang,^{46,h} S. S. Jiang,³⁹ T. J. Jiang,¹⁶ X. S. Jiang,^{1,58,63} Y. Jiang,⁶³ J. B. Jiao,⁵⁰ Z. Jiao,²³ S. Jin,⁴² Y. Jin,⁶⁶ M. Q. Jing,^{1,63} T. Johansson,⁷⁵ S. Kabana,³³ N. Kalantar-Nayestanaki,⁶⁴ X. L. Kang,⁹ X. S. Kang,⁴⁰ R. Kappert,⁶⁴ M. Kavatsyuk,⁶⁴ B. C. Ke,⁸⁰ A. Khoukaz,⁶⁸ R. Kiuchi,¹ R. Kliemt,¹³ L. Koch,³⁷ O. B. Kolcu,^{62a} B. Kopf,³ M. K. Kuessner,³ X. Kui,¹ A. Kupsc,^{44,75} W. Kühn,³⁷ J. J. Lane,⁶⁷ J. S. Lange,³⁷ P. Larin,¹⁸ A. Lavania,²⁶ L. Lavezzi,^{74a,74c} T. T. Lei,^{71,i} Z. H. Lei,^{71,58} H. Leithoff,³⁵ M. Lellmann,³⁵ T. Lenz,³⁵ C. Li,⁴³ C. Li,⁴⁷ C. H. Li,³⁹ Cheng Li,^{71,58} D. M. Li,⁸⁰ F. Li,^{1,58} G. Li,¹ H. Li,^{71,58} H. B. Li,^{1,63} H. J. Li,¹⁹ H. N. Li,^{56,j} Hui Li,⁴³ J. R. Li,⁶¹ J. S. Li,⁵⁹ J. W. Li,⁵⁰ Ke Li,¹ L. J. Li,^{1,63} L. K. Li,¹ Lei Li,⁴⁸ M. H. Li,⁴³ P. R. Li,^{38,k,l} S. X. Li,¹² T. Li,⁵⁰ W. D. Li,^{1,63} W. G. Li,^{1,a} X. H. Li,^{71,58} X. L. Li,⁵⁰ Xiaoyu Li,^{1,63} Y. G. Li,^{46,h} Z. J. Li,⁵⁹ Z. X. Li,¹⁵ Z. Y. Li,⁵⁹ C. Liang,⁴² H. Liang,^{71,58} H. Liang,^{1,63} H. Liang,³⁴ Y. F. Liang,⁵⁴ Y. T. Liang,^{31,63} G. R. Liao,¹⁴ L. Z. Liao,⁵⁰ J. Libby,²⁶ A. Limphirat,⁶⁰ D. X. Lin,^{31,63} T. Lin,¹ B. J. Liu,¹ B. X. Liu,⁷⁶ C. Liu,³⁴ C. X. Liu,¹ D. Liu,^{18,71} F. H. Liu,⁵³ Fang Liu,¹ Feng Liu,⁶ G. M. Liu,^{56,j} H. Liu,^{38,k,l} H. B. Liu,¹⁵ H. M. Liu,^{1,63} Huanhuan Liu,¹ Huihui Liu,²¹ J. B. Liu,^{71,58} J. L. Liu,⁷² J. Y. Liu,^{1,63} K. Liu,¹ K. Y. Liu,⁴⁰ Ke Liu,²² L. Liu,^{71,58} L. C. Liu,⁴³ Lu Liu,⁴³ M. H. Liu,^{12,g} P. L. Liu,¹ Q. Liu,⁶³ S. B. Liu,^{71,58} T. Liu,^{12,g} W. K. Liu,⁴³ W. M. Liu,^{71,58} X. Liu,^{38,k,l} Y. Liu,^{38,k,l} Y. B. Liu,⁴³ Z. A. Liu,^{1,58,63} Z. Q. Liu,⁵⁰ X. C. Lou,^{1,58,63} F. X. Lu,⁵⁹ H. J. Lu,²³ J. G. Lu,^{1,58} X. L. Lu,¹ Y. Lu,⁷ Y. P. Lu,^{1,58} Z. H. Lu,^{1,63} C. L. Luo,⁴¹ M. X. Luo,⁷⁹ T. Luo,^{12,g} X. L. Luo,^{1,58} X. R. Lyu,⁶³ Y. F. Lyu,⁴³ F. C. Ma,⁴⁰ H. L. Ma,¹ J. L. Ma,^{1,63} L. L. Ma,⁵⁰ M. M. Ma,^{1,63} Q. M. Ma,¹ R. Q. Ma,^{1,63} R. T. Ma,⁶³ X. Y. Ma,^{1,58} Y. Ma,^{46,h} Y. M. Ma,³¹ F. E. Maas,¹⁸ M. Maggiora,^{74a,74c} S. Maldaner,³ S. Malde,⁶⁹ A. Mangoni,^{28b} Y. J. Mao,^{46,h} Z. P. Mao,¹ S. Marcello,^{74a,74c} Z. X. Meng,⁶⁶ J. G. Messchendorp,^{13,64} G. Mezzadri,^{29a} H. Miao,^{1,63} T. J. Min,⁴² R. E. Mitchell,²⁷ X. H. Mo,^{1,58,63} N. Yu. Muchnoi,^{4,c} Y. Nefedov,³⁶ F. Nerling,^{18,e} I. B. Nikolaev,^{4,c} Z. Ning,^{1,58} S. Nisar,^{11,m} Y. Niu,⁵⁰ S. L. Olsen,⁶³ Q. Ouyang,^{1,58,63} S. Pacetti,^{28b,28c} X. Pan,⁵⁵ Y. Pan,⁵⁷ A. Pathak,³⁴ P. Patteri,^{28a} Y. P. Pei,^{71,58} M. Pelizaeus,³ H. P. Peng,^{71,58} K. Peters,^{13,e} J. L. Ping,⁴¹ R. G. Ping,^{1,63} S. Plura,³⁵ S. Pogodin,³⁶ V. Prasad,³³ F. Z. Qi,¹ H. Qi,^{71,58} H. R. Qi,⁶¹ M. Qi,⁴² T. Y. Qi,^{12,g} S. Qian,^{1,58} W. B. Qian,⁶³ C. F. Qiao,⁶³ J. J. Qin,⁷² L. Q. Qin,¹⁴ X. P. Qin,^{12,g} X. S. Qin,⁵⁰ Z. H. Qin,^{1,58} J. F. Qiu,¹ S. Q. Qu,⁶¹ C. F. Redmer,³⁵ K. J. Ren,³⁹ A. Rivetti,^{74c} V. Rodin,⁶⁴ M. Rolo,^{74c} G. Rong,^{1,63} Ch. Rosner,¹⁸ S. N. Ruan,⁴³ N. Salone,⁴⁴ A. Sarantsev,^{36,d} Y. Schelhaas,³⁵ K. Schoenning,⁷⁵ M. Scodeggio,^{29a,29b} K. Y. Shan,^{12,g} W. Shan,²⁴ X. Y. Shan,^{71,58} J. F. Shanguan,⁵⁵ L. G. Shao,^{1,63} M. Shao,^{71,58} C. P. Shen,^{12,g} H. F. Shen,^{1,63} W. H. Shen,⁶³ X. Y. Shen,^{1,63} B. A. Shi,⁶³ H. C. Shi,^{71,58} J. L. Shi,¹² J. Y. Shi,¹ Q. Q. Shi,⁵⁵ R. S. Shi,^{1,63} X. Shi,^{1,58} J. J. Song,¹⁹ T. Z. Song,⁵⁹ W. M. Song,^{34,1} Y. J. Song,¹² Y. X. Song,^{46,h,n} S. Sosio,^{74a,74c} S. Spataro,^{74a,74c} F. Stielor,³⁵ Y. J. Su,⁶³ G. B. Sun,⁷⁶ G. X. Sun,¹ H. Sun,⁶³ H. K. Sun,¹ J. F. Sun,¹⁹ K. Sun,⁶¹ L. Sun,⁷⁶ S. S. Sun,^{1,63} T. Sun,^{1,63} W. Y. Sun,³⁴ Y. Sun,⁹ Y. J. Sun,^{71,58} Y. Z. Sun,¹ Z. T. Sun,⁵⁰ Y. X. Tan,^{71,58} C. J. Tang,⁵⁴ G. Y. Tang,¹ J. Tang,⁵⁹ Y. A. Tang,⁷⁶ L. Y. Tao,⁷² Q. T. Tao,^{25,i} M. Tat,⁶⁹ J. X. Teng,^{71,58} V. Thoren,⁷⁵ W. H. Tian,⁵⁹ W. H. Tian,⁵² Y. Tian,^{31,63} Z. F. Tian,⁷⁶ I. Uman,^{62b} B. Wang,¹ B. L. Wang,⁶³ Bo Wang,^{71,58} C. W. Wang,⁴² D. Y. Wang,^{46,h} F. Wang,⁷² H. J. Wang,^{38,k,l} H. P. Wang,^{1,63} K. Wang,^{1,58} L. L. Wang,¹ M. Wang,⁵⁰ Meng Wang,^{1,63} S. Wang,^{12,g} S. Wang,^{38,k,l} T. Wang,^{12,g} T. J. Wang,⁴³ W. Wang,⁵⁹ W. Wang,⁷² W. H. Wang,⁷⁶ W. P. Wang,^{71,58,o} X. Wang,^{46,h} X. F. Wang,^{38,k,l} X. J. Wang,³⁹ X. L. Wang,^{12,g} Y. Wang,⁶¹ Y. D. Wang,⁴⁵ Y. F. Wang,^{1,58,63} Y. H. Wang,⁴⁷ Y. N. Wang,⁴⁵ Y. Q. Wang,¹ Yaqian Wang,^{17,1} Yi Wang,⁶¹ Z. Wang,^{1,58} Z. L. Wang,⁷² Z. Y. Wang,^{1,63} Ziyi Wang,⁶³ D. Wei,⁷⁰ D. H. Wei,¹⁴ F. Weidner,⁶⁸ S. P. Wen,¹ C. W. Wenzel,³ U. W. Wiedner,³ G. Wilkinson,⁶⁹ M. Wolke,⁷⁵ L. Wollenberg,³ C. Wu,³⁹ J. F. Wu,^{1,63} L. H. Wu,¹ L. J. Wu,^{1,63} X. Wu,^{12,g} X. H. Wu,³⁴ Y. Wu,⁷¹ Y. J. Wu,³¹ Z. Wu,^{1,58}

L. Xia,^{71,58} X. M. Xian,³⁹ T. Xiang,^{46,h} D. Xiao,^{38,k,l} G. Y. Xiao,⁴² H. Xiao,^{12,g} S. Y. Xiao,¹ Y. L. Xiao,^{12,g} Z. J. Xiao,⁴¹
 C. Xie,⁴² X. H. Xie,^{46,h} Y. Xie,⁵⁰ Y. G. Xie,^{1,58} Y. H. Xie,⁶ Z. P. Xie,^{71,58} T. Y. Xing,^{1,63} C. F. Xu,^{1,63} C. J. Xu,⁵⁹ G. F. Xu,¹
 H. Y. Xu,^{66,p} Q. J. Xu,¹⁶ Q. N. Xu,³⁰ W. Xu,^{1,63} W. L. Xu,⁶⁶ X. P. Xu,⁵⁵ Y. C. Xu,⁷⁷ Z. P. Xu,⁴² Z. S. Xu,⁶³ F. Yan,^{12,g}
 L. Yan,^{12,g} W. B. Yan,^{71,58} W. C. Yan,⁸⁰ X. Q. Yan,^{1,63} H. J. Yang,^{51,f} H. L. Yang,³⁴ H. X. Yang,¹ Tao Yang,¹ Y. Yang,^{12,g}
 Y. F. Yang,⁴³ Y. X. Yang,^{1,63} Yifan Yang,^{1,63} Z. W. Yang,^{38,k,l} M. Ye,^{1,58} M. H. Ye,⁸ J. H. Yin,¹ Z. Y. You,⁵⁹ B. X. Yu,^{1,58,63}
 C. X. Yu,⁴³ G. Yu,^{1,63} T. Yu,⁷² X. D. Yu,^{46,h} C. Z. Yuan,^{1,63} L. Yuan,² S. C. Yuan,¹ X. Q. Yuan,¹ Y. Yuan,^{1,63} Z. Y. Yuan,⁵⁹
 C. X. Yue,³⁹ A. A. Zafar,⁷³ F. R. Zeng,⁵⁰ X. Zeng,^{12,g} Y. Zeng,^{25,i} Y. J. Zeng,^{1,63} X. Y. Zhai,³⁴ Y. H. Zhan,⁵⁹ A. Q. Zhang,^{1,63}
 B. L. Zhang,^{1,63} B. X. Zhang,¹ D. H. Zhang,⁴³ G. Y. Zhang,¹⁹ H. Zhang,⁷¹ H. H. Zhang,³⁴ H. H. Zhang,⁵⁹ H. Q. Zhang,^{1,58,63}
 H. Y. Zhang,^{1,58} J. J. Zhang,⁵² J. L. Zhang,²⁰ J. Q. Zhang,⁴¹ J. W. Zhang,^{1,58,63} J. X. Zhang,^{38,k,l} J. Y. Zhang,¹ J. Z. Zhang,^{1,63}
 Jianyu Zhang,⁶³ Jiawei Zhang,^{1,63} L. M. Zhang,⁶¹ L. Q. Zhang,⁵⁹ Lei Zhang,⁴² P. Zhang,¹ Q. Y. Zhang,^{39,80}
 Shuihan Zhang,^{1,63} Shulei Zhang,^{25,i} X. D. Zhang,⁴⁵ X. M. Zhang,¹ X. Y. Zhang,⁵⁰ X. Y. Zhang,⁵⁵ Y. Zhang,⁶⁹ Y. Zhang,⁷²
 Y. T. Zhang,⁸⁰ Y. H. Zhang,^{1,58} Yan Zhang,^{71,58} Yao Zhang,¹ Z. H. Zhang,¹ Z. L. Zhang,³⁴ Z. Y. Zhang,⁴³ Z. Y. Zhang,⁷⁶
 G. Zhao,¹ J. Zhao,³⁹ J. Y. Zhao,^{1,63} J. Z. Zhao,^{1,58} Lei Zhao,^{71,58} Ling Zhao,¹ M. G. Zhao,⁴³ S. J. Zhao,⁸⁰ Y. B. Zhao,^{1,58}
 Y. X. Zhao,^{31,63} Z. G. Zhao,^{71,58} A. Zhemchugov,^{36,b} B. Zheng,⁷² J. P. Zheng,^{1,58} W. J. Zheng,^{1,63} Y. H. Zheng,⁶³ B. Zhong,⁴¹
 X. Zhong,⁵⁹ H. Zhou,⁵⁰ L. P. Zhou,^{1,63} X. Zhou,⁷⁶ X. K. Zhou,⁶ X. R. Zhou,^{71,58} X. Y. Zhou,³⁹ Y. Z. Zhou,^{12,g} J. Zhu,⁴³
 K. Zhu,¹ K. J. Zhu,^{1,58,63} L. Zhu,³⁴ L. X. Zhu,⁶³ S. H. Zhu,⁷⁰ S. Q. Zhu,⁴² T. J. Zhu,^{12,g} W. J. Zhu,^{12,g} Y. C. Zhu,^{71,58}
 Z. A. Zhu,^{1,63} J. H. Zou,¹ and J. Zu^{71,58}

(BESIII Collaboration)

¹*Institute of High Energy Physics, Beijing 100049, People's Republic of China*

²*Beihang University, Beijing 100191, People's Republic of China*

³*Bochum Ruhr-University, D-44780 Bochum, Germany*

⁴*Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*

⁵*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

⁶*Central China Normal University, Wuhan 430079, People's Republic of China*

⁷*Central South University, Changsha 410083, People's Republic of China*

⁸*China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*

⁹*China University of Geosciences, Wuhan 430074, People's Republic of China*

¹⁰*Chung-Ang University, Seoul, 06974, Republic of Korea*

¹¹*COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*

¹²*Fudan University, Shanghai 200433, People's Republic of China*

¹³*GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*

¹⁴*Guangxi Normal University, Guilin 541004, People's Republic of China*

¹⁵*Guangxi University, Nanning 530004, People's Republic of China*

¹⁶*Hangzhou Normal University, Hangzhou 310036, People's Republic of China*

¹⁷*Hebei University, Baoding 071002, People's Republic of China*

¹⁸*Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*

¹⁹*Henan Normal University, Xinxiang 453007, People's Republic of China*

²⁰*Henan University, Kaifeng 475004, People's Republic of China*

²¹*Henan University of Science and Technology, Luoyang 471003, People's Republic of China*

²²*Henan University of Technology, Zhengzhou 450001, People's Republic of China*

²³*Huangshan College, Huangshan 245000, People's Republic of China*

²⁴*Hunan Normal University, Changsha 410081, People's Republic of China*

²⁵*Hunan University, Changsha 410082, People's Republic of China*

²⁶*Indian Institute of Technology Madras, Chennai 600036, India*

²⁷*Indiana University, Bloomington, Indiana 47405, USA*

^{28a}*INFN Laboratori Nazionali di Frascati, I-00044, Frascati, Italy*

^{28b}*INFN Sezione di Perugia, I-06100, Perugia, Italy*

^{28c}*University of Perugia, I-06100, Perugia, Italy*

^{29a}*INFN Sezione di Ferrara, I-44122, Ferrara, Italy*

^{29b}*University of Ferrara, I-44122, Ferrara, Italy*

³⁰*Inner Mongolia University, Hohhot 010021, People's Republic of China*

³¹*Institute of Modern Physics, Lanzhou 730000, People's Republic of China*

³²*Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia*

- ³³*Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile*
- ³⁴*Jilin University, Changchun 130012, People's Republic of China*
- ³⁵*Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*
- ³⁶*Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia*
- ³⁷*Justus-Liebig-Universität Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany*
- ³⁸*Lanzhou University, Lanzhou 730000, People's Republic of China*
- ³⁹*Liaoning Normal University, Dalian 116029, People's Republic of China*
- ⁴⁰*Liaoning University, Shenyang 110036, People's Republic of China*
- ⁴¹*Nanjing Normal University, Nanjing 210023, People's Republic of China*
- ⁴²*Nanjing University, Nanjing 210093, People's Republic of China*
- ⁴³*Nankai University, Tianjin 300071, People's Republic of China*
- ⁴⁴*National Centre for Nuclear Research, Warsaw 02-093, Poland*
- ⁴⁵*North China Electric Power University, Beijing 102206, People's Republic of China*
- ⁴⁶*Peking University, Beijing 100871, People's Republic of China*
- ⁴⁷*Qufu Normal University, Qufu 273165, People's Republic of China*
- ⁴⁸*Renmin University of China, Beijing 100872, People's Republic of China*
- ⁴⁹*Shandong Normal University, Jinan 250014, People's Republic of China*
- ⁵⁰*Shandong University, Jinan 250100, People's Republic of China*
- ⁵¹*Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China*
- ⁵²*Shanxi Normal University, Linfen 041004, People's Republic of China*
- ⁵³*Shanxi University, Taiyuan 030006, People's Republic of China*
- ⁵⁴*Sichuan University, Chengdu 610064, People's Republic of China*
- ⁵⁵*Soochow University, Suzhou 215006, People's Republic of China*
- ⁵⁶*South China Normal University, Guangzhou 510006, People's Republic of China*
- ⁵⁷*Southeast University, Nanjing 211100, People's Republic of China*
- ⁵⁸*State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China*
- ⁵⁹*Sun Yat-Sen University, Guangzhou 510275, People's Republic of China*
- ⁶⁰*Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand*
- ⁶¹*Tsinghua University, Beijing 100084, People's Republic of China*
- ^{62a}*Turkish Accelerator Center Particle Factory Group, Istinye University, 34010, Istanbul, Turkey*
- ^{62b}*Near East University, Nicosia, North Cyprus, 99138, Mersin 10, Turkey*
- ⁶³*University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China*
- ⁶⁴*University of Groningen, NL-9747 AA Groningen, The Netherlands*
- ⁶⁵*University of Hawaii, Honolulu, Hawaii 96822, USA*
- ⁶⁶*University of Jinan, Jinan 250022, People's Republic of China*
- ⁶⁷*University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom*
- ⁶⁸*University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany*
- ⁶⁹*University of Oxford, Keble Road, Oxford OX13RH, United Kingdom*
- ⁷⁰*University of Science and Technology Liaoning, Anshan 114051, People's Republic of China*
- ⁷¹*University of Science and Technology of China, Hefei 230026, People's Republic of China*
- ⁷²*University of South China, Hengyang 421001, People's Republic of China*
- ⁷³*University of the Punjab, Lahore-54590, Pakistan*
- ^{74a}*University of Turin and INFN, University of Turin, I-10125, Turin, Italy*
- ^{74b}*University of Eastern Piedmont, I-15121, Alessandria, Italy*
- ^{74c}*INFN, I-10125, Turin, Italy*
- ⁷⁵*Uppsala University, Box 516, SE-75120 Uppsala, Sweden*
- ⁷⁶*Wuhan University, Wuhan 430072, People's Republic of China*
- ⁷⁷*Yantai University, Yantai 264005, People's Republic of China*
- ⁷⁸*Yunnan University, Kunming 650500, People's Republic of China*
- ⁷⁹*Zhejiang University, Hangzhou 310027, People's Republic of China*
- ⁸⁰*Zhengzhou University, Zhengzhou 450001, People's Republic of China*

^aDeceased.^bAlso at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.^cAlso at the Novosibirsk State University, Novosibirsk, 630090, Russia.^dAlso at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia.^eAlso at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

^fAlso at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China.

^gAlso at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China.

^hAlso at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China.

ⁱAlso at School of Physics and Electronics, Hunan University, Changsha 410082, China.

^jAlso at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China.

^kAlso at MOE Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People's Republic of China.

^lAlso at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China.

^mAlso at the Department of Mathematical Sciences, IBA, Karachi 75270, Pakistan.

ⁿAlso at Ecole Polytechnique Federale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland.

^oAlso at Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany.

^pAlso at School of Physics, Beihang University, Beijing 100191, China.